



Improving Antidepressant Adherence Rates in an Outpatient Behavioral Setting: A Quality Improvement Project

TAIWO ABIOYE, PMHNP-BC

DNPV 767 – PROJECT

TOURO UNIVERSITY NEVADA

PROJECT MEMBER(S): DR. JOHNSTON, DR. D'ERRICO

JUNE 22, 2021

ACKNOWLEDGEMENT

I want to extend my appreciation to my academic mentor Dr. Heidi Johnston; her unwavering support, guidance, patience, and commitment to my success have been overwhelming. I am also thankful to Dr. D'errico for her guidance and direction through this journey. Special appreciation to my project mentor Dr. Charmeka Lipscomb; your support is appreciated. I would also like to show appreciation to this great school's entire staff, especially my instructors Dr. Grimm, Dr. Carrion, Dr. Zabriskie, and Dr. Thompson, for their assistance and feedback through this program.



INTRODUCTION

- Depression is a mood disorder that causes persistent sadness and affects the quality of human life and psychosocial functioning.
- The project aims to evaluate antidepressant adherence in patients with depression by implementing an evidence-based guideline and protocol by mental health providers.
- This DNP project proved to improve the adherence of patients through the implementation of an antidepressant adherence protocol.
- The quality improvement DNP project contributes significantly improved provider's knowledge about depression and increased antidepressant adherence in patients with depression.



BACKGROUND

- The National Institute of Mental health reported depression as the most prevalent mental health disorder in the United States of America, with over 16 million people clinically diagnosed with depression.
- The national recommendations for managing depression are effective management and treatment with therapeutic support, pharmacotherapy, and psychotherapy.
- Nonadherence to antidepressants is the failure to adhere to medical provider's treatment recommendations, which can negatively impact health and cause a significant challenge to patient outcomes.
- Nonadherence to antidepressants leads to low quality of life, poor health outcomes, increased risks of treatment relapse, vulnerability to suicide, and increased costs to the patient and health care system.
- Studies show that 60% of nonadherence to antidepressants was higher in minority groups, and they are less likely to refill their antidepressants, and that 60% of patients discontinue antidepressants use within three months.



PROJECT PURPOSE

- The purpose of this DNP project is:
 - The purpose of the DNP project was to implement a quality improvement project to improve antidepressant adherence in adult patients with depression and enhance the knowledge of the mental health providers of an outpatient behavioral setting.

PROJECT PROBLEM

This quality improvement (QI) project problem is the lack of evidence-based guidelines to improve antidepressant adherence in patients with depression.

PROJECT QUESTION



Will implementing an evidence-based guideline on antidepressant adherence improve the adherence rate in adult patients of 18 -75 years within four weeks of the project Implementation phase?

PROJECT OBJECTIVES



1. Develop and implement evidenced-based toolkit and guidelines for providers prescribing antidepressants to enhance antidepressant medication adherence.
2. Educate providers and present the developed evidenced-based protocol to providers, evaluate their understanding of the guidelines.
3. Evaluate their understanding of the guidelines and assess improved knowledge with a pre-and post-knowledge tests.
4. Show a 5% increase in antidepressant medication adherence within four weeks of implementation.
5. Improve early intervention rates for patients at risk for nonadherence to antidepressants/


LITERATURE REVIEW

- Implementation of evidence-based guidelines to improve antidepressant adherence rates in an outpatient behavioral setting.
- Impact of the problem; Antidepressant nonadherence can lead to decreased quality of life and poor health outcomes for patients.
- Addressing the Problem with Current Evidence identified factors influencing nonadherence in patients on antidepressants, including stigma, fear of the medication's adverse reaction, the ineffectiveness of antidepressants, and disbelief about diagnosis and severity of depression.

TABLE 2. BARRIERS TO ANTIDEPRESSANT ADHERENCE

Care access
Concern about potential sexual dysfunction, weight gain, and other adverse effects
Fear of antidepressants
Lack of an adherence strategy
Medication cost
Misattributions about the causes of depression
Negative attitudes and a perceived stigma
Perception of low symptom severity
Perception that depression is a character flaw
Unrealistic expectations

THEORETICAL MODEL



Kurt Lewin's Change Theory

1. Recognize the need for change
2. Determine what needs to change
3. Encourage the replacement of old behaviors and attitudes
4. Ensure there is strong support from management
5. Manage and understand the doubts and concerns

1. Plan the changes
2. Implement the changes
3. Help employees to learn new concept or points of view

1. Changes are reinforced and stabilized
2. Integrate changes into the normal way of doing things
3. Develop ways to sustain the change
4. Celebrate success

PROJECT DESIGN



- The design for this project utilized a quality improvement (QI) design focused on implementing the best practice guidelines that outline the process for providers and nurses to follow, to improve antidepressant adherence rates in adult patients.
- The DNP quality improvement (QI) project's primary goal is to maintain the implementation of the best practice guideline in order to enhance adherence in patients on antidepressants.



PROJECT SETTING

The setting is an outpatient behavioral clinic located in El Paso Texas. This clinic provides medication management, psychosocial and rehabilitative programs, low energy and neurofeedback systems, and individual and family counseling to patients aged 18-75 years. The clinic serves approximately 3,000 patients and sees more than 190 patients daily, with approximately 80% Hispanics, 10% Caucasians, 5% African Americans, and 5% Middle Eastern and Asians.

PROJECT PLAN

- The DNP project was implemented during a four-week time frame.
- The implementation phase included:
 - Implementing the intervention
 - Data collection
 - Evaluating the project results



IMPLEMENTATION

- Week 1:** The project lead will meet with the providers, give a pretest to the medical providers, and provide education, followed by a post-test.
- Week 2:** Weekly retrospective chart reviews and implementation of the evidenced-based toolkit that outlines the process for providers (Nurse Practitioners, Physician Assistants, and Physicians) to follow in enhancing adherence to antidepressant in adult patients at the practice site will continue. Meeting with the providers/stakeholders about updates on the project. Weekly meetings with the providers as needed to assess adherence to the protocol, check on the progress, and outcomes of the implementation.
- Week 3:** Implementation of the evidenced-based toolkit that outlines the process for providers to follow in enhancing adherence to antidepressant adherence in adult patients at the practice site will continue. Random chart reviews post implementation of the protocol
- Week 4:** Assess patient's adherence to antidepressant medications, final data collection and analysis of data began.

DATA COLLECTION

- This is a Quality improvement (QI) project.
- The QI project did not require the Institutional Review Board (IRB).
- Approval received from the CEO, Chief medical director, and stakeholders of the project site.
- Direct recruitment of population of interest and utilized in-person and zoom meetings.
- Pre and post test of provider knowledge administered electronically during the educational intervention sessions.
- Chart audit and data collection performed over a four- week period.
- No personal data collected.
- Data was exported to IBM SPSS Statistics.


RESULTS

- The result of the pre-test and post-test were compared, and the increase in score between the pre and post-test showed a knowledge increase evidenced by a mean score of 81% versus a score of 99% on the post-test.
- Descriptively, post-test (adherence section), all providers increased their adherence to protocol with an overall rate of 77% from 0%.
- Significant differences between pretest and posttest scores for knowledge.
- Pre-test scores ranged from 60-90% and post-test scores ranged from 95-100%.
- Post-test- scores improved in comparison to the pre-intervention scores 60% to over 95%.




RESULTS

Provider Pre-Post Antidepressant Adherence Knowledge Test



RESULTS

Provider Pre-Post Protocol Adherence



RESULTS

- The t-paired test was performed to assess if providers knowledge score post-test was different from the pre-test knowledge score.
- The binomial test was used to assess if the protocol adherence by providers pre-implementation is statistically significant post-implementation.
- The chart audit pre- and post-implementation of provider teaching and antidepressant protocol use showed a significant improvement in new patients on antidepressants.
- There was a significant difference between patient pre-post adherence to antidepressants after the project implementation ($p=0.017$).
- Pre-implementation audit of 30 patient's antidepressant adherence rate prior to the teaching was at 60%, and 80% post-implementation representing a 20% increase.
- Descriptively, post-test (adherence section), all providers increased their adherence to protocol with an overall rate of 77% from 0%.

Table 1.

Pre and Post Pre-Post Antidepressant Adherence Knowledge Test Paired Samples Statistics

Paired Samples Statistics					
Pair 1	pre	Mean 81.00	N 6	Std. Deviation 2.000	Std. Error Mean .816
	post	99.00	6	2.000	.816

Table 2.

Pre and Post Pre-Post Antidepressant Adherence Knowledge Test Paired Samples Correlations

Paired Samples Correlations				
Pair		N	Correlation	Sig.
Pair 1	pre & post	6	.250	.633

Table 3.

Pre and Post Pre-Post Antidepressant Adherence Knowledge Test Paired Samples Test

Paired Samples Test				
Pair 1	pre - post	t	df	Sig. (2-tailed)
		-18.000	5	.000

RESULTS

Table 4.

Pre and Post Pre-Post Antidepressant Adherence Knowledge Test Paired Differences

Paired Samples Test

		Paired Differences				
		Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference	
					Lower	Upper
Pair 1	pre - post	-18.000	2.449	1.000	-20.571	-15.429

Pre and Post Pre-Post Antidepressant Adherence Knowledge Test Paired Samples Effect Sizes

Paired Samples Effect Sizes						
			Standardizer ^a	Point Estimate	95% Confidence Interval	
					Lower	Upper
Pair 1	pre - post	Cohen's d	2.449	-7.348	-11.835	-2.909
		Hedges' correction	2.655	-6.781	-10.920	-2.684

LIMITATIONS


- Difficulties resulting from social distancing and associated complexities of the COVID-19 pandemic restrictions, the clinical site changed services to an online/virtual platform.
- Educational training via zoom Vs in person.
- Short timeline for implementing the project intervention, which took place over four weeks.
- Chart audits on antidepressant adherence was collected for four weeks after the educational training of the providers was implemented.
- The number of provider participants at the practice site for this DNP project was limited.
- There were ten prescribing medical providers at the facility initially, however, only five prescribing medical providers were working at the practice site at the time of the project implementation.
- Minimal sample size, smaller than the anticipated number of providers for this project.

SUSTAINABILITY



- Stakeholders such as the Chief Medical Officer and Chief Operations Officer, provided support needed to sustain the goals of this quality improvement project.
- Integration of the protocol for all prescribers at the project site.
- An in-service training will be provided to educate newly hired medical providers and nursing staff on the antidepressant adherence protocol.
- The medical director has given the go-ahead to allow this intervention to serve as a healthcare model for the clinic.

EVALUATION



- Evaluation entailed an assessment of provider knowledge increase regarding the importance of antidepressant adherence.
- It evaluated the provider's knowledge and awareness of antidepressant adherence protocol, thereby determining an increase in the percentage of adherence in patients on antidepressants compared to pre-implementation.
- The change in adherence rates pre-and post-implementation determined was analyzed using paired t-test analysis.
- A binomial test was used to analyze the results of the pre-and post-protocol adherence by providers.
- The chart audit was completed, the data was compiled into an Excel spreadsheet, and the analysis was done with the SPSS software.

FUTURE DISSEMINATION

- The project will be prepared as a PowerPoint presentation and delivered to the academic staff and students of Touro University on June 2021 in Las Vegas, NV.
- The project will also be submitted to the DNP repository.
- The DNP candidate will the DNP QI project has presented the results of this project to the stakeholders and providers in the current healthcare organizations.
- The project lead also plans to disseminate the findings of this project to the Texas chapter of Black Nurses Rock Professional Organization.
- It is also planned to be disseminated to the DNPs of color (DOP) professional conference on July 22, 2021.
- An abstract will be submitted for presentation at the 2021 AANP Fall Conference in Hollywood, Florida, dated September 23, 2021 – September 26, 2021.
- The project will also be disseminated through an abstract submission for publication and presentation to the Global Alliance and the International Society of Psychiatric-Mental Health Nurses (ISPN) dated March 15-20, 2022, at Redondo Beach, CA.

CONCLUSION

- Increased knowledge of the mental health providers.
- Utilization of antidepressant protocol showed improvement in adherence to antidepressants in patients with depression.
- Mental health provider's attitudes towards the protocol were improved by education.
- Utilizing the adherence protocol helped enhance favorable outcomes for the patients.





Thank You Everyone

Taiwo Abioye



**ANY
QUESTIONS**

REFERENCES

- American Psychiatric Association. (2015). Diagnostic and statistical manual of mental disorders (5th Ed.). American Psychiatric Publishing.
- Herzog, D. P., Wagner, S., Ruckes, C., Tadic, A., Roll, S. C., Härter, M., & Lieb, K. (2017). Guideline adherence of antidepressant treatment in outpatients with major depressive disorder: a naturalistic study. *European Archives of Psychiatry and Clinical Neuroscience*, 267(8), 711-721. <http://dx.doi.org/10.1007/s00406-017-0798-6>
- Henein, F., Prabhakar, D., Peterson, E. L., Williams, L. K., & Ahmedani, B. K. (2016). A prospective study of antidepressant adherence and suicidal ideation among adults. *Primary Care Companion for CNS Disorders*, 18(6) DOI:10.4088/PCC.16101935
- Cipriani, A., Furukawa, T. A., Salanti, G., Chaimani, A., Atkinson, L. Z., Ogawa, Y., Geddes, J. R. (2018). Comparative efficacy and acceptability of 21 antidepressant drugs for the acute treatment of adults with major depressive disorder: A systematic review and network meta-analysis. *The Lancet (British Edition)*, 391(10128), 1357-1366. DOI:10.1016/s0140-6736(17)32802-7
- Costa, E., Pecorelli, S., Giardini, A., Savin, M., Menditto, E., Lehane, E., Laosa O., Monaco, A., Marengoni, A. (2015). Interventional tools to improve medication adherence: Review of the literature. *Patient Preference and Adherence*, 9, 1303. doi:10.2147/ppa.s87551
- Srimongkon, P., Aslani, P., & Chen, T. (2018). Consumer-related factors influencing antidepressant adherence in unipolar depression: A qualitative study. *Patient Preference and Adherence*, 12, 1863-1873. DOI:10.2147/ppa.s160728